

A Unified Model of Short and Long Gamma-Ray Bursts, X-Ray Rich Gamma-Ray Bursts, and X-Ray Flashes

Ryo Yamazaki

Department of Earth and Space Science, Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan

Kunihito Ioka

Physics Department and Center for Gravitational Wave Physics, 104 Davey Laboratory, Pennsylvania State University, University Park, PA 16802, USA

Takashi Nakamura, Kenji Toma

Theoretical Astrophysics Group Department of Physics, Kyoto University Kyoto 606-8502, Japan

Abstract

We propose a possible unified model of short and long gamma-ray bursts (GRBs), X-ray rich GRBs, and X-ray flashes. The jet of a GRB is assumed to consist of multiple sub-jets or sub-shells (i.e., an inhomogeneous jet model). The multiplicity of the sub-jets along a line of sight n_s is an important parameter. If n_s is large ($\gg 1$) the event looks like a long GRB, while if $n_s = 1$, the event looks like a short GRB. Finally, when $n_s = 0$, the event looks like an X-ray flash or an X-ray rich GRB. Furthermore, our model may also explain the bimodal distributions of T_{90} duration of BATSE-GRBs.

Key words: gamma-ray bursts — gamma rays: theory

1 Introduction

While the association with the supernova is almost established or strongly suggested for the long gamma-ray bursts (GRBs) (Stanek et al., 2003; Hjorth et al.,

Email address: ryo@vega.ess.sci.osaka-u.ac.jp (Ryo Yamazaki).

2003), the origins of short GRBs and X-ray flashes (XRFs) remain unclear. The observed event rate of short GRBs is about a third of the long GRBs while the observed event rate of XRFs is also about a third. Although there may be a possible bias effect to these statistics, these numbers are, in an astrophysical sense, the same or comparable. If these phenomena arise from essentially different origins, the similar number of events is just by chance. On the other hand, if they are related, the similar number of events is natural and the ratio of the event rate tells us something about the geometry of the central engine. We propose a unified model in which the central engine of short GRBs, long GRBs and XRFs is the same and the apparent differences come essentially from different viewing angles. For details, see Yamazaki et al. (2004b) and Toma et al. (2004).

2 Unified Model

Roughly speaking, short GRBs are similar to the first 1 sec of long GRBs (Ghirlanda et al., 2004), which suggests that the difference between short and long GRBs is just the number of pulses, and each pulse is essentially the same. Thus, we may consider that each pulse is produced by essentially the same unit or the sub-jet, and the GRB jet consists of many sub-jets. If many sub-jets point to our line of sight, the event looks like the long GRB while if a single sub-jet points to us, the event looks like a short GRB. Since we can observe only the angular size of $\sim \gamma^{-1}$ within the GRB jet with the Lorentz factor γ , different observers will see different number of sub-jets depending on the distribution of sub-jets within the GRB jet.

XRFs also appear to be related to GRBs. Softer and dimmer GRBs smoothly extend to the XRFs (Heise et al., 2001; Kippen et al., 2002; Sakamoto et al., 2004). Other properties of the XRFs are also similar to those of the GRBs, suggesting that XRFs are in fact soft and dim GRBs. In the sub-jet model, XRFs are naturally expected when our line of sight is off-axis to any sub-jets (Nakamura, 2000; Ioka & Nakamura, 2001; Yamazaki et al., 2002, 2003b,c, 2004a).

In the following, we show a numerical simulation to demonstrate how the event looks so different depending on the viewing angle in our unified model (Yamazaki et al., 2004b). The whole jet, with an opening half-angle of $\Delta\theta_{\text{tot}} \sim 0.2$ rad, consists of 350 sub-jets. All sub-jets are assumed to have the same intrinsic properties, e.g., the opening half-angle $\Delta\theta_{\text{sub}}^{(j)} = 0.02$ rad, and so on. We consider the case in which the angular distribution of sub-jets is given by $P(\vartheta^{(j)}) \propto \exp[-(\vartheta^{(j)}/\vartheta_c)^2/2]$, where we adopt $\vartheta_c = 0.1$ rad (Zhang et al., 2004). In this case, sub-jets are concentrated on the central region. For our adopted parameters, isolated sub-jets exist near the edge of the whole jet with

the multiplicity $n_s = 1$ and there exists a viewing angle where no sub-jets are launched. The left, middle, and right panels of Fig. 1 show the angular distributions of sub-jets and the directions of three selected lines of sight, the observed light curves in the γ -ray bands, and the observed time-integrated spectra, respectively. Note here in the left panel of Fig. 1, “A” represents the center of the whole jet and is hidden by the lines of sub-jets.

Long GRB: When we observe the source from the $\vartheta = 0$ axis (case “A”), we see spiky temporal structures (the upper-middle panel of Fig. 1) and $E_p \sim 300$ keV which are typical for the long GRBs. We may identify case “A” as long GRBs.

XRF and X-ray rich GRB: When the line of sight is away from any sub-jets (case “B”), soft and dim prompt emission, i.e. XRFs or X-ray rich GRBs are observed with $E_p = 10 \sim 20$ keV and ~ 4 orders of magnitude smaller fluence than that of case “A” (the right panel of Fig. 1). The burst duration is comparable to that in case “A”. These are quite similar to the characteristics of XRFs. We may identify the case “B” as XRFs or X-ray rich GRBs.

Short GRB: If the line of sight is inside an isolated sub-jet (case “C”), its observed pulse duration is ~ 50 times smaller than case “A”. Contributions to the observed light curve from the other sub-jets are negligible, so that the fluence is about a hundredth of the case “A”. These are quite similar to the characteristics of short GRBs. However the hardness ratio ($= S(100 - 300 \text{ keV})/S(50 - 100 \text{ keV})$) is about 3 which is smaller than the mean hardness of short GRBs (~ 6). Ghirlanda et al. (2004) suggested that the hardness of short GRBs is due to the large low-energy photon index $\alpha_B \sim -0.58$ so that if the central engine launches $\alpha_B \sim -0.58$ sub-jets to the periphery of the core where n_s is small, we may identify the case “C” as the short-hard GRBs. In other words, the hardness of 3 comes from $\alpha_B = -1$ in our simulation so that if $\alpha_B \sim -0.58$, the hardness will be 6 or so. We suggest here that not only the isotropic energy but also the photon index may depend on ϑ . Another possibility is that if short GRBs are the first 1 sec of the activity of the central engine, the spectrum in the early time might be $\alpha_B \sim -0.58$ for both the sub-jets in the core and the envelope. This is consistent with a high KS test probability for E_p and α_B (Ghirlanda et al., 2004). These possibilities may have something to do with the origin of $\alpha_B \sim -1$ for the long GRBs.

3 Discussions

Let $\Delta\theta_{\text{sub}}$, ϑ_c and $\bar{n}_s$ be the typical opening half-angle of the sub-jet, the core size of the whole jet and the mean multiplicity in the core. Then the total number of the sub-jets (N_{tot}) is estimated as $N_{\text{tot}} = \bar{n}_s(\vartheta_c/\Delta\theta_{\text{sub}})^2 \sim 10^3$, so that the total energy of each sub-jet is $\sim 10^{48}$ erg. In our model, the event




Fig. 1. (Left) The angular distribution of 350 sub-jets confined in the whole GRB jet in our simulation. The whole jet has the opening half-angle of $\Delta\theta_{\text{tot}} = 0.2$ rad. The sub-jets have the same properties. The axes and the angular size of sub-jets are represented by crosses and the dotted circles, respectively. “A” represents the center of the whole jet and is hidden by the lines of sub-jets. (Middle) The observed γ -ray light curves, corresponding the cases “A” and “C” in the left panel. (Right) Time-integrated energy spectrum of the emission from the multiple sub-jets for the observers denoted by “A”, “B”, and “C”. The source is located at $z = 1$.

rate of long GRBs is in proportion to ϑ_c^2 . Let M be the number of sub-jets in the envelope of the core with a multiplicity $n_s = 1$. Then the event rate of short GRBs is in proportion to $M\Delta\theta_{\text{sub}}^2$, so that $M \sim 10$ is enough to explain the event rate of short GRBs.

Since the core may be regarded as a uniform jet, our model for the XRFs is analogous to the off-axis uniform jet model (Yamazaki et al., 2002, 2003b, 2004a). However the afterglow could have a different behavior between the core-envelope sub-jet model and the uniform jet model. In the uniform jet model, the afterglows of XRFs should resemble the orphan afterglows that initially have a rising light curve (Yamazaki et al., 2003a; Granot et al., 2002). An orphan afterglow may be actually observed in XRF 030723 (Fynbo et al., 2004), but the light curve may peak too early (Zhang et al., 2004). The optical afterglow of XRF 020903 is not observed initially (< 0.9 days) but may not be consistent with the orphan afterglow (Soderberg et al., 2004). These problems could be overcome by introducing a Gaussian tail with a high Lorentz factor around the uniform jet (Zhang et al., 2004) since the energy redistribution effects may bring the rising light curve to earlier times (Zhang et al., 2004; Kumar & Granot, 2003). The afterglow of a short GRB is difficult to predict since it could resemble both the orphan and normal afterglow depending on the sub-jet configuration within the envelope.

It is also found that our model can reproduce the bimodal distribution of T_{90} duration of GRBs observed by BATSE (Toma et al., 2004). In our model, The duration of $n_s = 1$ burst is determined by the angular spreading time of one sub-jet emission, while that of $n_s \geq 2$ burst is determined by the time interval between the observed first pulse and the last one. These two different time scales naturally lead a division of the burst T_{90} durations into the short and

long ones. It has commonly been said that the observed bimodal distribution of T_{90} durations of BATSE bursts shows the different origins of short and long GRBs. However, the bimodal distribution is also available as a natural consequence of our unified model of short and long GRBs. The clear prediction of our unified model is that short GRBs should be associated with energetic SNe. Indeed, one of short GRBs shows the possible association with a SN (Germany et al., 2000). Even if the SNe are not identified with short GRBs due to some observational reasons we predict that the spatial distribution of short GRBs in host galaxies should be similar to that of the long GRBs. Another prediction is that short GRBs should have the same total kinetic energies as long GRBs, which might be confirmed by radio calorimetry.

Interestingly, our model has predicted short XRFs or short X-ray rich GRBs (Yamazaki et al., 2004b). They are observed when isolated sub-jets are viewed slightly off-axis. The observed short XRF 040924 may be a kind of these bursts (Fenimore et al., 2004).

References

- Fenimore, E. et al. 2004, GCN Circ. 2735
 Fynbo, J. P. U., Sollerman, J., Hjorth, J. et al., 2004, ApJ, 609, 962
 Ghirlanda, G., Ghisellini, G., & Celotti, A., 2004, A&A, 422, L55
 Germany, L. M., Reiss, D. J., Sadler, E. M. et al., 2000, ApJ, 533, 320
 Granot, J., Panaitescu, A., Kumar, P., & Woosley, S. E., 2002, ApJ, 570, L61
 Heise, J., in 't Zand, J., Kippen, R. M., & Woods, P. M., 2001, in Proc. Second Rome Workshop: Gamma-Ray Bursts in the Afterglow Era, ed. E. Costa, F. Frontera, & J. Hjorth (Berlin: Springer), 16
 Hjorth, J., Sollerman, J., Möller, P. et al., 2003, Nature, 423, 847
 Ioka, K., & Nakamura, T., 2001, ApJ, 554, L163
 Kippen, R. M., Woods, P. M., Heise, J. et al. 2002, AIP Conf. Proc. 662, 244 (astro-ph/0203114)
 Kumar, P., & Granot, J., 2003, ApJ, 591, 1075
 Nakamura, T., 2000, ApJ, 534, L159
 Sakamoto, T., Lamb, D. Q., Graziani, C. et al., 2004, ApJ, 602, 875
 Soderberg, A. M., Kulkarni, S. R., Berger, E. et al., 2004, ApJ, 606, 994
 Stanek, K. Z. Matheson, T., Garnavich, P. M. et al., 2003, ApJ, 591, L17
 Toma, K., Yamazaki, R., & Nakamura, T., 2004, astro-ph/0407012
 Yamazaki, R., Ioka, K., & Nakamura, T., 2002, ApJ, 571, L31
 Yamazaki, R., Ioka, K., & Nakamura, T., 2003a, ApJ, 591, 283
 Yamazaki, R., Ioka, K., & Nakamura, T., 2003b, ApJ, 593, 941
 Yamazaki, R., Yonetoku, D., & Nakamura, T., 2003c, ApJ, 594, L79
 Yamazaki, R., Ioka, K., & Nakamura, T., 2004a, ApJ, 606, L33
 Yamazaki, R., Ioka, K., & Nakamura, T., 2004b, ApJ, 607, L103

Zhang, B., Dai, X., Lloyd-Ronning, N. M., & Mészáros, P., 2004, ApJ, 601,
L119